

WHEAT Receival Standards - Soft & Stockfeed 2021/2022



These Standards apply to Wheat Receivals for ALL Clients delivering into Emerald Grain and are to be applied on individual truck loads and must not be averaged over a number of loads

QUALITY PARAMETER		SFE1	SFE2	SFW1	SFWR				
GTA Standards CSG Number		140	143	151	152				
CATEGORY									
VARIETY	Varietal Restrictions Apply	VR	Yes	Yes	Yes	Yes			
PROTEIN	Min %	PR	N/A	N/A	N/A	N/A			
	Max %	PR	9.5	10.5	N/A	N/A			
MOISTURE	Max %	MO	12.5	12.5	12.5	12.5			
TEST WEIGHT	Min kg/hl	TW	76.0	76.0	70.0	70.0			
SCREENINGS (Below Screen)	The total material passing through the 2.0mm Screen after 40 shakes	SC	8.0	8.0	10.0	10.0			
Unmillable Material Above the Screen	Whiteheads (Grain Removed) Chaff Backbone, Wild Radish Pods, Milk Thistle, Skeleton Weed Flowers and other seed pods not otherwise listed	SA	0.6	0.6	1.2	1.2			
Defective Grains (% by count per 300 grain sample unless otherwise stated)									
Sprouted	Max % by Count	SP	Nil	Nil	N/A	N/A			
Falling Number	Min Seconds	FN	300	300	N/A	N/A			
The NIL Tolerance of Sprouted Grain does not apply if a Falling Number analysis is Conducted									
Stained	Max % by Count	FS	10.0	15.0	15.0	15.0			
Pink Stained	Max % by Count	PF	2.0	5.0	5.0	5.0			
White Grain Disorder/Head Scab	Max % by Count	WG	1.0	1.0	1.0	1.0			
Field Fungi	Max by Count per ½ Litre	FF	10	10	10	10			
Dry Green or Sappy	Max % by Count	GS	1.0	5.0	10.0	10.0			
Distorted	Max % by Count	FR	1.0	5.0	10.0	10.0			
Severely Damaged - Above 2.0mm Screen	Max by Count per ½ Litre	BB	1	1	1	1			
All Smuts except Loose Smut	Entire Load	SM	Nil	Nil	Nil	Nil			
Insect Damaged	Max % by Count	ID	1.0	1.0	2.0	2.0			
Foreign Seed Contaminants – Max (count total per half litre unless otherwise stated)									
Type 1:	Colocynthis, Double Gees / Spiny Emex / Three Cornered Jack, Field Poppy, Long Head Poppy, Mexican Poppy, New Zealand Spinach, Opium Poppy, Jute, Wild Poppy, Horned Poppy	S1	8	8	8	8			
Type 2:	Caster Oil Plant, Coriander, Crow Garlic / Wild Garlic, Darling Pea, Peanut seeds and pods, Ragweed, Rattlepods, Starburr, St Johns Wort, Parthenium Weed	S2	Nil	Nil	Nil	Nil			
Type 3a:	Bathurst Burr, Bellvine, Branched Broomrape, Bulls Head / Caltrop / Cats Heads, Cape Tulip, Cottonseed, Dodder, Noogoora Burr, Thornapple	3a	2	2	2	2			
Type 3b:	Vetch (Tare), Vetch (Commercial)	3b	4	4	4	4			
Type 3c:	Heliotrope (Blue), Heliotrope (Common)	3c	2 Pods 8 Seeds	2 Pods 8 Seeds	2 Pods 8 Seeds	2 Pods 8 Seeds			
Type 4:	Field Bindweed, Cutleaf Mignonette seeds, Darnel (Drake Seed), **Hexham Scent/Melilot (King Island), Hoary Cress, Paddy Melon, Variegated Thistle, Nightshades, Skeleton Weed, Mintweed **Hexham Scent is only acceptable if no tainting odour is present	S4	20	20	20	20			
Type 5:	Creeping Knapweed / Russian Knapweed, Sesbania Pea, Patterson's Curse / Salvation Jane	S5	40	40	40	40			
Type 6:	Columbus Grass, Johnson Grass, Saffron Thistle	S6	10	10	50	50			
Type 7a:	Adzuki Beans, Broad Bean, Chickpeas, Corn (Maize), Onion Weed pods, Cowpea, Faba Beans, Lentils, Lupin, Safflower, Soybean, Pea (Field), Medic Pods, Sunflower and any other seed or pod greater than 5 mm in diameter	7a	1	1	10	10			
Type 7b:	2 Row Barley, 6 Row Barley, Australian Bindweed, Black Bindweed, Brome Grass, Cereal Rye, Durum, Carrot Weed, Red Wheat, Oats (Wild / Black / Common / Sand), Rice, Ryegrass on Stalk, Sorghum (grain), Triticale, Turnip Weed, and any other foreign seeds not specified in types 1-7, SFS or SA	7b	50	50	150	150			
Small Foreign Seeds:	All Foreign Seeds not specified in Types 1-7(b) that fall below the 2.0mm screen during the Screenings process	SS	0.6	0.6	1.2	1.2			
Other Contaminants									
Pickling Compounds	Entire Load	PI	Nil	Nil	Nil	Nil			
Chemicals Not Approved for Wheat	Entire Load	CH	Nil	Nil	Nil	Nil			
Ryegrass Ergot – Total Length of all pieces laid end to end	Length in cm per ½ litre	RE	2.0	2.0	2.0	2.0			
Cereal Ergot	Max Count per ½ litre	CE	1	1	1	1			
Stored Grain Insects & Pea Weevil – Live	Entire Load	LGI	Nil	Nil	Nil	Nil			
Insects – Large, dead or alive	Max Count per ½ litre	LI	3	3	3	3			
Insects – Small, dead or alive – Included Dead Stored Grain Insects	Max Count per ½ litre	SI	10	10	10	10			
Earcokle	Max Count per ½ litre	EC	10	10	15	15			
Snails – dead or alive	Max Count per ½ litre	SN	1	1	10	10			
Loose Smut	Max Count per ½ litre	LS	3	3	3	3			
Sand – 0.06-2.00mm in size, includes Earth and Stones below 2.00mm Screen	Max Count per ½ litre	ES	20	20	50	50			
Earth – Above 2.00mm Screen, Max 5mm in Diameter	Max Count per ½ litre	EA	1	1	3	3			
Stones – Above 2.00mm Screen (weight in grams)	Max Weight per 2.5litres	SE	4.0	4.0	4.0	4.0			
Other Foreign Material	Max % by Weight	OF	0.1	0.1	0.1	0.1			
Objectionable Material	Entire Load	OM	Nil	Nil	Nil	Nil			
Sticks	A Stick is defined as ligneous material with dimensions of greater than 3cm in length and/or 1cm in diameter Smaller material is classified as Screenings or Unmillable material above the screen								
Odour or Tainting Agents	Any commercially unacceptable odour is defined as a sour or musty odour imparted on the wheat that is not considered natural which may or may not be physically detectable through the presence of foreign material. Included but is not limited to plant parts and seed of Eucalyptus spp.								
Restricted Animal Material	Presence of meat meal, bone meal, blood meal, poultry offal meal or any other animal proteins Restricted material also included carcasses or dead animals such as rats and mice								
Other	Animal excreta, glass, concrete, metal, fertilizer								
Water	The addition of water to grain prior to delivery is a prohibited practice								